

Cohomology of hyperkähler manifolds via arithmetic harmonic analysis

Tamás Hausel

Royal Society URF at University of Oxford

University of Texas at Austin

<http://www.math.utexas.edu/~hausel/talks.html>

November 2005

Geometry Seminar, Kyoto University

Representation theory and hyperkähler varieties

- **[Grojnowski 1996, Nakajima 1997]** Representation of the Heisenberg algebra on $\bigoplus_{n=1}^{\infty} H^*((\mathbb{C}^2)^{[n]})$ the cohomology of Hilbert schemes of n points on \mathbb{C}^2
- **[Nakajima 1995,1998]** Representations of a Kac-Moody algebra on $\bigoplus_{\mathbf{v},\mathbf{w}} H^*(\mathcal{M}(\mathbf{v},\mathbf{w}))$ the cohomology of Nakajima quiver varieties
- **[Nakajima 2001]** Representations of quantum affine algebras on $\bigoplus_{\mathbf{v},\mathbf{w}} K^*(\mathcal{M}(\mathbf{v},\mathbf{w}))$ the K -theory of Nakajima quiver varieties
- **[Haiman 2002]** Representation theory of S_n in $K^*((\mathbb{C}^2)^{[n]})$ the K -theory of Hilbert scheme of n points on $\mathbb{C}^2 \implies$ Macdonald polynomials
- **[Beilinson, Drinfeld ~1995-]** Geometric Langlands programme \implies representation theory of $G(F_C)$, a reductive group over the function field of a complex curve, in the non-commutative geometry of hyperkähler moduli spaces in the non-abelian Hodge theory of C for G^L .

Motivation: mirror symmetry

- A pair of n dimensional Calabi-Yau manifolds (X, Y) satisfy the topological mirror test if

$$H^{p,q}(X) = H^{n-p,q}(Y)$$

- A pair of n dimensional Calabi-Yau manifolds (X, Y) are Strominger-Yau-Zaslow mirror pairs if they map to the same real n -dimensional manifold B , so that the generic fibers are dual special Lagrangian tori

Diffeomorphic spaces in the non-Abelian Hodge theory of a genus g curve C :

$$\mathcal{M}_{\text{Dol}}^d(GL(n, \mathbb{C})) := \left\{ \begin{array}{l} \text{moduli space of semistable rank } n \\ \text{degree } d \text{ Higgs bundles on } C \end{array} \right\}$$

$$\mathcal{M}_{\text{DR}}^d(GL(n, \mathbb{C})) := \left\{ \begin{array}{l} \text{moduli space of flat } GL(n, \mathbb{C})\text{-connections} \\ \text{on } C \setminus \{p\}, \text{ with holonomy } e^{\frac{2\pi id}{n}} Id \text{ around } p \end{array} \right\}$$

$$\mathcal{M}_{\text{B}}^d(GL(n, \mathbb{C})) := \{A_1, B_1, \dots, A_g, B_g \in GL(n, \mathbb{C}) \mid A_1^{-1} B_1^{-1} A_1 B_1 \dots A_g^{-1} B_g^{-1} A_g B_g = \xi_n Id\} / GL(n, \mathbb{C})$$

Theorem 1 (Hausel–Thaddeus 2003). *In the following diagram*

$$\begin{array}{ccc}
 \mathcal{M}_{\text{Dol}}^d(PGL(n)) & \longrightarrow & \mathcal{M}_{\text{Dol}}^d(SL(n)) \\
 \downarrow \chi_{PGL(n)} & & \downarrow \chi_{SL(n)} \\
 \mathcal{H}_{PGL(n)} & \cong & \mathcal{H}_{SL(n)}.
 \end{array}$$

the generic fibers of the Hitchin maps $\chi_{PGL(n)}$ and $\chi_{SL(n)}$ are dual Abelian varieties.

\Downarrow

$\mathcal{M}_{\text{DR}}^d(PGL(n))$ and $\mathcal{M}_{\text{DR}}^d(SL(n))$ satisfy the SYZ construction for a pair of mirror symmetric Calabi-Yau manifolds.

Topological Mirror Test

Conjecture 1 (Hausel–Thaddeus 2003). *For all $d, e \in \mathbb{Z}$, satisfying $(d, n) = (e, n) = 1$, we have*

$$E_{\text{st}}^{B^e}(x, y; \mathcal{M}_{\text{DR}}^d(SL(n, \mathbb{C}))) = E_{\text{st}}^{\hat{B}^d}(x, y; \mathcal{M}_{\text{DR}}^e(PGL(n, \mathbb{C}))).$$

Conjecture 2 (Hausel–R-Villegas 2004).

$$E_{\text{st}}^{B^e}(x, y, \mathcal{M}_{\text{B}}^d(SL(n, \mathbb{C}))) = E_{\text{st}}^{\hat{B}^d}(x, y, \mathcal{M}_{\text{B}}^e(PGL(n, \mathbb{C}))).$$

Holomorphic symplectic quotients

- \mathbb{V} finite dimensional complex vector space, G complex reductive group
- representation $\rho : G \rightarrow GL(\mathbb{V})$ of G on \mathbb{V} , inducing $\varrho : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathbb{V})$.
- inducing an action of G on $\mathbb{V} \times \mathbb{V}^*$, preserving the natural symplectic structure, with moment map: $\mu : \mathbb{V} \times \mathbb{V}^* \rightarrow \mathfrak{g}^*$ defined by $\mu(v, w)(X) = \langle \varrho(X)v, w \rangle$
- for $\xi \in (\mathfrak{g}^*)^G$ the holomorphic symplectic quotient: $\mathbb{V} \times \mathbb{V}^* //_{\xi} G := (\mu^{-1}(\xi)) // G$ carries a natural hyperkähler metric at its smooth points
- Examples: affine toric hyperkähler varieties when G abelian; Nakajima's quiver varieties, when ρ is constructed from a quiver, e.g. semisimple adjoint orbits in $\mathfrak{gl}(n, \mathbb{C})$

Group-valued symplectic quotients

$G = GL(n)$; $C = \mathbb{P}^1$ is a genus 0 curve with generic semisimple conjugacy classes $\tilde{C}_1, \dots, \tilde{C}_k \subset GL(n, \mathbb{C})$ at punctures $p_1, \dots, p_k \in C$. The *character variety* is

$$\mathcal{M}_B = \{(A_1, \dots, A_k) \mid A_i \in \tilde{C}_i, A_1 \dots A_k = Id\} // G(\mathbb{C})$$

Riemann-Hilbert monodromy map
 \cong

$$\mathcal{M}_{DR} := \left\{ \begin{array}{l} \text{moduli space of flat} \\ GL(n, \mathbb{C})\text{-connections on } C \setminus \{p_1, \dots, p_k\} \\ \text{with holonomy around } p_i \text{ lying in } \tilde{C}_i \end{array} \right\}$$

Mixed Hodge Structure of Deligne

- $\bigoplus_{p,q} H^{p,q;k}(M)$ is the associated graded to the weight and Hodge filtrations on the cohomology $H^k(M, \mathbb{C})$ of a complex algebraic variety M
- $h^{p,q;k} = \dim(H^{p,q;k}(M))$, the *mixed Hodge numbers*
- $H(M; x, y, t) = \sum_{p,q,k} h^{p,q;k}(M) x^p y^q t^k$, the *mixed Hodge polynomial*
- $P(M; t) = H(M; 1, 1, t)$, the *Poincaré polynomial*
- $E(M; x, y) = x^n y^n H(1/x, 1/y, -1)$, the *E-polynomial* of a smooth variety M .

Arithmetic and topological content of the E-polynomial

Theorem 2 (... , Ito 2004, Katz 2005). *If M is a smooth quasi-projective variety defined over \mathbb{Z} and*

$$\#\{M(\mathbb{F}_q)\} = E(q)$$

is a polynomial in q , then

$$E(M; x, y) = E(xy).$$

- MHS on $H^*(M, \mathbb{C})$ is *pure* if $h^{p,q;k} = 0$ unless $p + q = k \Leftrightarrow H(M; x, y, t) = (xyt^2)^n E(\frac{-1}{xt}, \frac{-1}{yt}) \Rightarrow P(M; t) = H(M; 1, 1, t) = t^{2n} E(\frac{-1}{t}, \frac{-1}{t})$; examples of varieties with pure MHS: smooth projective varieties, \mathcal{M}_{Dol} , \mathcal{M}_{DR} , Nakajima's quiver varieties
- in general the pure part of $H(M; x, y, t)$ is $PH(M; x, y) = \text{Coeff}_{T^0} \left(H(M; xT, yT, tT^{-1}) \right)$; which, for a smooth M , is always the image of the cohomology of a smooth compactification

Fourier Transform on finite groups

- Γ finite group; the convolution of the functions $f_1, \dots, f_k : \Gamma \rightarrow \mathbb{C}$ is defined as

$$f_1 \star f_2 \star \dots \star f_k(h) =$$

$$|\Gamma|^{(1-k)/2} \sum_{\substack{g_1, g_2, \dots, g_k \in \Gamma \\ g_1 g_2 \dots g_k = h}} f_1(g_1) f_2(g_2) \dots f_k(g_k)$$

- Fourier transform of function $f : \Gamma \rightarrow \mathbb{C}$ at $\rho : \Gamma \rightarrow \text{End}(V_\rho)$ irrep.:

$$\mathcal{F}(f, \rho) = |\Gamma|^{-1/2} \sum_{g \in \Gamma} f(g) \rho(g) \in \text{End}(V_\rho),$$

- $\mathcal{F}(f_1 \star f_2, \rho) = \mathcal{F}(f_1, \rho) \circ \mathcal{F}(f_2, \rho)$

- Fourier inversion formula:

$$|\Gamma|^{-1/2} \sum_{\rho \in \text{Irr}(\Gamma)} \dim(V_\rho) \text{tr} \left(\mathcal{F}(f, \rho) \circ \rho(h^{-1}) \right) = f(h)$$

Fourier Transform for \mathcal{M}_B

Setup:

- $G = GL(n)$
- $C = \mathbb{P}^1$, with punctures $a_1, \dots, a_k \in \mathbb{P}^1$
- $\tilde{C}_i \subset GL(n)$ fixed semisimple conjugacy classes
- $\mathcal{M}_B = \{(A_1, A_2, \dots, A_k) \mid A_i \in \tilde{C}_i, A_1 \cdots A_k = 1\} // G(\mathbb{C})$

Theorem 3 (Frobenius 1896, Hausel–Villegas 2004).

$$\#\{\mathcal{M}_B(\mathbb{F}_q)\} = \frac{|Z(G(\mathbb{F}_q))|}{|G(\mathbb{F}_q)|} 1_{\tilde{C}_1} \star \cdots \star 1_{\tilde{C}_k}(1) = \sum_{\chi \in \text{Irr}(G(\mathbb{F}_q))} \frac{\chi(1)^2 |Z(G(\mathbb{F}_q))|}{|G(\mathbb{F}_q)|^2} \prod_i \frac{\chi(\tilde{C}_i(\mathbb{F}_q))}{\chi(1)} |\tilde{C}_i(\mathbb{F}_q)|$$

is a polynomial $E(q)$ in $q \Rightarrow$

$$E(\mathcal{M}_B, x, y) = E(xy)$$

Example Assume $n = 3$, and all the conjugacy classes \tilde{C}_i are regular semisimple:

$$\begin{aligned}
 E(\mathcal{M}_B; q) = & \\
 & \frac{\left((q+1)(q^2+q+1)\right)^k}{(q^3-1)^2(q^2-1)^2} - \frac{\left(3q^2(q+1)\right)^k}{q^4(q^2-1)^2(q-1)^2} \\
 & + 1/3 \frac{\left(6q^3\right)^k}{q^6(q-1)^4} + \frac{\left(2q^2(q^2+q+1)\right)^k}{q^4(q^3-1)^2(q-1)^2} \\
 & + \frac{\left(q^3(q+1)(q^2+q+1)\right)^k}{q^6(q^3-1)^2(q^2-1)^2} - \frac{\left(3q^3(q+1)\right)^k}{q^6(q^2-1)^2(q-1)^2}.
 \end{aligned}$$

Conjecture 3 (Hausel 2004). When $n = 3$, \tilde{C}_i are regular semisimple, $h_{N-j}^{i-j} = h_{N+j}^{i+j}$ for

$$\begin{aligned}
H(\mathcal{M}_B, q, t) = & \sum h_j^i q^j t^i = \\
& \frac{\left((qt^2 + 1) (q^2 t^4 + qt^2 + 1) \right)^k}{(q^3 t^6 - 1) (q^3 t^4 - 1) (q^2 t^4 - 1) (q^2 t^2 - 1)} \\
& - \frac{\left(3 q^2 t^4 (qt^2 + 1) \right)^k}{q^4 t^8 (q^2 t^4 - 1) (q^2 t^2 - 1) (qt^2 - 1) (q - 1)} \\
& + \frac{1}{3} \frac{6^k (qt^2)^{3k}}{q^6 t^{12} (qt^2 - 1)^2 (q - 1)^2} \\
& + \frac{\left(q^2 t^4 (2 q^2 t^2 + qt^2 + q + 2) \right)^k}{q^4 t^8 (q^3 t^4 - 1) (q^3 t^2 - 1) (qt^2 - 1) (q - 1)} \\
& + \frac{\left(q^3 t^6 (q + 1) (q^2 + q + 1) \right)^k}{q^6 t^{12} (q^3 t^2 - 1) (q^3 - 1) (q^2 t^2 - 1) (q^2 - 1)} \\
& - \frac{\left(3 q^3 t^6 (q + 1) \right)^k}{q^6 t^{12} (q^2 t^2 - 1) (q^2 - 1) (qt^2 - 1) (q - 1)},
\end{aligned}$$

The Riemann-Hilbert map

- $\mathfrak{g} = \mathfrak{gl}(n)$
- $C = \mathbb{P}^1$ with punctures $a_1, \dots, a_k \in \mathbb{P}^1$
- \mathcal{C}_i semisimple adjoint orbit in $\mathfrak{g}(\mathbb{C})$
- $Q = \{(A_1, \dots, A_k) \mid A_i \in \mathcal{C}_i, A_1 + \dots + A_k = 0\} // G(\mathbb{C})$, Nakajima's star-shaped quiver variety
- Q is smooth when \mathcal{C}_i are generic
- " $Q \subset \mathcal{M}_{\text{DR}}$ ", a point in Q gives the meromorphic flat $GL(n, \mathbb{C})$ -connection $\sum A_i \frac{dz}{z-a_i}$ on the trivial bundle on C .
- $\tilde{\mathcal{C}}_i = \exp(2\pi i \mathcal{C}_i) \subset G(\mathbb{C})$ is the corresponding conjugacy class
- the *Riemann-Hilbert monodromy map*

$$\nu_a : Q \rightarrow \mathcal{M}_{\text{B}}$$

is given by sending the flat connection to its holonomy.

The purity conjecture

Conjecture 4. *If C_i are generic, then*

$$\nu_a^* : PH^*(\mathcal{M}_B) \xrightarrow{\cong} H^*(Q)$$

Example

- $n = 3, k = 3, C_i$ regular semisimple
- Q is \cong to an E_6 ALE space,
- $\mathcal{M}_B \cong \mathcal{M}_{\text{Dol}}$ an elliptic fibration with singular fibre of type \hat{E}_6 .
- $P_t(Q) = 1 + 6t^2$
- $H(\mathcal{M}_B; q, t) = 1 + 6qt^2 + q^2t^2$

↓

Conjecture is true in this case

Fourier transform for Q

- $\Psi : \mathbb{F}_q \rightarrow \mathbb{C}$ non-trivial additive character
- $f : \mathfrak{g}(\mathbb{F}_q) \rightarrow \mathbb{C}$ its Fourier transform $\mathcal{F}(f) : \mathfrak{g}^*(\mathbb{F}_q) \rightarrow \mathbb{C}$ at a $Y \in \mathfrak{g}^*(\mathbb{F}_q)$

$$\mathcal{F}(f)(Y) := |\mathfrak{g}(\mathbb{F}_q)|^{-1/2} \sum_{X \in \mathfrak{g}(\mathbb{F}_q)} f(X) \Psi(\langle X, Y \rangle).$$

- $\delta_{\mathcal{C}_i} : \mathfrak{g}(\mathbb{F}_q) \rightarrow \mathbb{C}$ characteristic function of $\mathcal{C}_i \subset \mathfrak{g}(\mathbb{F}_q)$

Theorem 4 (Hausel–Villegas 2004).

$$\#\{Q(\mathbb{F}_q)\} = \frac{|Z(G(\mathbb{F}_q))| |\mathfrak{g}(\mathbb{F}_q)|^{\frac{k-2}{2}}}{|G(\mathbb{F}_q)| \sum_{X \in \mathfrak{g}(\mathbb{F}_q)} \mathcal{F}(\delta_{\mathcal{C}_1})(X) \cdots \mathcal{F}(\delta_{\mathcal{C}_k})(X)}$$

Theorem 5. *Letellier's character table for $\mathfrak{gl}(3, \mathbb{F}_q)$ implies that when $n = 3$ and all adjoint orbits C_i are regular semi-simple*

$$\begin{aligned}
 P(Q; t) = & \frac{\left((t^2 + 1) (t^4 + t^2 + 1) \right)^k}{(t^6 - 1) (t^4 - 1)} \\
 & - \frac{\left(3t^4 (t^2 + 1) \right)^k}{t^8 (t^4 - 1) (t^2 - 1)} + 1/3 \frac{6^k (t^2)^{3k}}{t^{12} (t^2 - 1)^2} \\
 & - \frac{\left(t^4 (t^2 + 2) \right)^k}{t^8 (t^2 - 1)} + t^{6k-12} + \frac{(3t^6)^k}{t^{12} (t^2 - 1)}.
 \end{aligned}$$

Conjecture 5 (Hausel–Letellier–Villegas 2005).

$g \geq 0, k > 0, \mu = \{\mu^1, \dots, \mu^k\} \in \mathcal{P}(n)^{\{1, \dots, k\}},$
 $X_1 = \{x_{1,1}, x_{1,2}, \dots\}, \dots, X_k = \{x_{k,1}, x_{k,2}, \dots\},$

$$V_\mu^n := H(\mathcal{M}_B(C_\mu); q, -t) \frac{(qt^2)^{(2-2g-k)\frac{n(n-1)}{2}} (1-qt)^{2g}}{(1-q)(1-qt^2)}$$

$$(qt^2)^{\sum_i^k n(\bar{\mu}^i)} m_{\mu^1}(X_1) \dots m_{\mu^k}(X_k),$$

$$Z_n(q, t) = \exp \left(\sum_{d=1}^{\infty} \sum_{\mu} \frac{1}{d} V_\mu^n(q^d, t^d, X_1^d, \dots, X_k^d) \right),$$

$$\mathcal{H}_\lambda^g(q, t) = \prod_{x \in d(\lambda)} \frac{(qt^2)^{(2-2g)l(x)} (1 - q^{h(x)} t^{2l(x)+1})^{2g}}{(1 - q^{h(x)} t^{2l(x)+2}) (1 - q^{h(x)} t^{2l(x)})}.$$

$$H_\lambda(X; q, t) = \sum_{|\rho|=|\lambda|} K_{\rho\lambda}(q, t) s_\rho(X),$$

Macdonald (q, t) -symmetric functions, $K_{\rho\lambda}(q, t)$ (q, t) -Kostka polynomials, $s_\rho(X)$ Schur and $m_\mu(X)$ monomial symmetric functions

$$\prod_{n=1}^{\infty} Z_n(q, t) = \sum_{\lambda \in \mathcal{P}} \left(\prod_{i=1}^k H_\lambda(X_i; q, qt^2) \right) \mathcal{H}_\lambda^g(q, t)$$

Theorem 6 (Hausel–Letellier–Villegas 2005).

Let $H_\mu(q, t) \stackrel{?}{=} H(\mathcal{M}_B(C_\mu), q, t)$ the conjectured mixed Hodge polynomial. Then

$$H_\mu(q, -1) = E(\mathcal{M}_B(C_\mu), q)$$

from the character table of $GL(n, \mathbb{F}_q)$ and the pure part

$$PH_\mu(t^2) = P(Q_\mu, t)$$

from the character table of $\mathfrak{gl}(n, \mathbb{F}_q)$.

Purity conjecture for $g > 0$

Setup

- C genus g curve, punctures $a_1, \dots, a_k \in C$
- $\tilde{C}_i \subset G$ semisimple conjugacy classes
- $\mathcal{M}_B = \{(A_1, B_1, \dots, A_g, B_g, C_1, \dots, C_k) \mid C_i \in \tilde{C}_i \mid A_1^{-1}B_1^{-1}A_1B_1 \dots A_g^{-1}B_g^{-1}A_gB_gC_1 \dots C_k = Id\} // G(\mathbb{C})$
- $\mathcal{C}_i \subset \mathfrak{g}$ semisimple adjoint orbit
- $Q = \{(A_1, B_1, \dots, A_g, B_g, C_1, \dots, C_k) \mid C_i \in \tilde{C}_i \mid A_1B_1 - B_1A_1 + \dots + A_gB_g - B_gA_g + C_1 + \dots + C_k = 0\} // G(\mathbb{C})$

Conjecture 6. *The pure part of the cohomology of \mathcal{M}_B is isomorphic with the cohomology of Q , in particular: $PH(\mathcal{M}_B; q, t) = P(Q; t)$*

Fourier transform for $(\mathbb{C}^2)^{[n]}$

- $g = 1$ and $k = 1$, and $\mathcal{C}_1 \subset GL(n)$ is a smallest non-central semisimple orbit

- $\mathcal{M}_B =$

$$\{(A_1, B_1, C) \mid C \in \mathcal{C}_1 \mid A_1^{-1} B_1^{-1} A_1 B_1 C = Id\} // GL(n)$$

- $Q \cong (\mathbb{C}^2)^{[n]}$

- Conjecture $\Rightarrow PH^*(\mathcal{M}_B) \cong H^*((\mathbb{C}^2)^{[n]}, \mathbb{C})$.

- [Nevins-Stafford 2003] \Rightarrow

$$\mathcal{M}_B \cong (\mathbb{C}^\times \times \mathbb{C}^\times)^{[n]}$$

Conjecture 7 (Hausel 2005). *In this case:*

$$\sum_{n=1}^{\infty} H(\mathcal{M}_B; q, t) T^n = \prod_{m=1}^{\infty} \frac{(1 + q^m t^{2m-1} T^m)^2}{(1 - q^{m-1} t^{2m-2} T^m)(1 - q^{m+1} t^{2m} T^m)}$$

Fourier transform for $\mathbb{V} \times \mathbb{V}^* // // \xi G$

- representation $\rho : G \rightarrow GL(\mathbb{V})$ of G on \mathbb{V} , inducing $\varrho : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathbb{V})$.
- inducing an action of G on $\mathbb{V} \times \mathbb{V}^*$, preserving the natural symplectic structure, with moment map: $\mu : \mathbb{V} \times \mathbb{V}^* \rightarrow \mathfrak{g}^*$ defined by $\mu(v, w)(X) = \langle \varrho(X)v, w \rangle$

Proposition 7 (Hausel 2005). *Let $\xi \in \mathfrak{g}(\mathbb{F}_q)^*$. The number of solutions of the equation $\mu(x) = \xi$ over the finite field \mathbb{F}_q is given by the formula:*

$$\begin{aligned} \#\{(v, w) \in \mathbb{V}(\mathbb{F}_q) \times \mathbb{V}(\mathbb{F}_q)^* \mid \mu(v, w) = \xi\} &= \\ |\mathfrak{g}(\mathbb{F}_q)|^{-1} |\mathbb{V}(\mathbb{F}_q)| \sum_{X \in \mathfrak{g}(\mathbb{F}_q)} C_\varrho(X) \frac{(q\delta_0(\langle X, \xi \rangle) - 1)}{q - 1} &= \\ &= |\mathfrak{g}(\mathbb{F}_q)|^{-1/2} |\mathbb{V}| \mathcal{F}(C_\varrho)(\xi), \end{aligned}$$

where $C_\varrho(X) = |\ker(\varrho(X))|$.

Theorem 8 (Hausel 2005). $Q = (\mathcal{V}, \mathcal{E})$ quiver, $\mathcal{V} = \{1, \dots, n\}$ vertices $\mathcal{E} \subset \mathcal{V} \times \mathcal{V}$ edges, $\mathbf{w} \in \mathbb{N}^{\mathcal{V}}$ dimension vector, $\mathcal{M}(\mathbf{v}, \mathbf{w})$ affine Nakajima quiver variety.

$$\begin{aligned} & \sum_{\mathbf{v} \in \mathbb{N}^{\mathcal{V}}} P_t(\mathcal{M}(\mathbf{v}, \mathbf{w})) t^{-d(\mathbf{v}, \mathbf{w})} T^{\mathbf{v}} = \\ & \sum_{\mathbf{v} \in \mathbb{N}^{\mathcal{V}}} T^{\mathbf{v}} \sum_{\lambda \in \mathcal{P}(\mathbf{v})} \frac{\left(\prod_{(i,j) \in \mathcal{E}} t^{-2n(\lambda^i, \lambda^j)} \right) \left(\prod_{i \in \mathcal{V}} t^{-2n(\lambda^i, (1^{\mathbf{w}_i}))} \right)}{\prod_{i \in \mathcal{V}} \left(t^{-2n(\lambda^i, \lambda^i)} \prod_k \prod_{j=1}^{m_k(\lambda^i)} (1-t^{2j}) \right)}, \\ & = \frac{\sum_{\mathbf{v} \in \mathbb{N}^{\mathcal{V}}} T^{\mathbf{v}} \sum_{\lambda \in \mathcal{P}(\mathbf{v})} \frac{\prod_{(i,j) \in \mathcal{E}} t^{-2n(\lambda^i, \lambda^j)}}{\prod_{i \in \mathcal{V}} \left(t^{-2n(\lambda^i, \lambda^i)} \prod_k \prod_{j=1}^{m_k(\lambda^i)} (1-t^{2j}) \right)}}{\prod_{i \in \mathcal{V}} \left(t^{-2n(\lambda^i, \lambda^i)} \prod_k \prod_{j=1}^{m_k(\lambda^i)} (1-t^{2j}) \right)}, \end{aligned}$$

where $d(\mathbf{v}, \mathbf{w}) = 2\sum_{(i,j) \in \mathcal{E}} \mathbf{v}_i \mathbf{v}_j + 2\sum_{i \in \mathcal{V}} \mathbf{v}_i (\mathbf{w}_i - \mathbf{v}_i)$ is the dimension of $\mathcal{M}(\mathbf{v}, \mathbf{w})$, $T^{\mathbf{v}} = \prod_{i \in \mathcal{V}} T_i^{\mathbf{v}_i}$ and $n(\lambda, \mu) = \sum_{i,j} \min(\lambda_i, \mu_j)$

Example $\mathcal{V} = \{1\}$, $\mathcal{E} = \{(1, 1)\}$, the twisted ADHM space of $U(k)$ Yang-Mills instantons of charge n on \mathbb{R}^4 is then $\mathcal{M}(n, k) = \mu^{-1}(Id_V) // G$, where

$$\mu(A, B, I, J) = [A, B] + IJ$$

is the ADHM equation. Theorem 7 then implies the result of Nakajima-Yoshioka 2004:

$$\sum_{n=0}^{\infty} P_t(\mathcal{M}(n, k)) T^n = \prod_{i=1}^{\infty} \prod_{b=1}^k \frac{1}{(1 - t^{2(k(i-1)+b-1)} T^i)}.$$